

Toward a Clean Sample of Ultra-Luminous X-ray Sources

M. López-Corredoira¹, C. M. Gutiérrez¹

¹ Instituto de Astrofísica de Canarias, C/Vía Láctea, s/n, E-38200 La Laguna (S/C de Tenerife), Spain

Received xxxx; accepted xxxx

ABSTRACT

Context. Observational follow-up programmes for the characterization of ultra-luminous X-ray sources (ULXs) require the construction of clean samples of such sources in which the contamination by foreground/background sources is minimum.

Aims. In this article we calculate the degree of foreground/background contaminants among the ULX sample candidates in the Colbert & Ptak (2002) catalogue and compare these computations with available spectroscopical identifications.

Methods. We use statistics based on known densities of X-ray sources and AGN/QSOs selected in the optical. The analysis is done individually for each parent galaxy. The existing identifications of the optical counterparts are compiled from the literature.

Results. More than a half of the ULXs, within twice the distance of the major axis of the 25 mag/arcsec² isophote from RC3 nearby galaxies and with X-ray luminosities $L_X[2-10 \text{ keV}] \geq 10^{39} \text{ erg/s}$, are expected to be high redshift background QSOs. A list of 25 objects (clean sample) confirmed to be real ULXs or to have a low probability of being contaminant foreground/background objects is provided.

Key words. Galaxies: statistics – quasars: general – X-rays: individuals: IXOs

1. Introduction

ULXs (ultraluminous X-ray sources) are objects associated with some nearby galaxies, and which emit in X-ray with luminosities above the Eddington limit (so they cannot be stellar-mass X-ray binaries), but which are well below the typical emission in the center of AGNs. The physical nature of ULXs still is unclear. One of the most interesting possibilities is that they are black holes of intermediate masses (IMBHs; Colbert & Mushotzky 1999; Makishima et al. 2000). Follow-up programmes over wide spectral ranges are essential for revealing the nature of these sources.

There is strong observational evidence that points to the existence of other kinds of objects linked to the parent galaxies: short-lived massive stars, usually in HII regions (e.g. Pakull & Mirioni 2002; Gao et al. 2003; Kaaret et al. 2004; Zampieri et al. 2004; Liu et al. 2004; Smith et al. 2005; Lehmann et al. 2005), tentatively associated with intermediate-mass black holes. The ULX phenomenon is indeed closely connected to star formation activities, since ULXs preferentially occur in spiral arms of late-type or starbursts rather than in early-type galaxies (Kilgard et al. 2002; Schwartz et al. 2004; Liu et al. 2005b). Actually, 70% of starburst galaxies host at least one ULX. However, as has been shown in the identification of several optical counterparts of ULXs (Lira et al. 2000; Arp et al. 2004; Gutiérrez & López-Corredoira 2005, 2006; Gutiérrez 2006; Wong et al. 2005; Clark et al. 2005; e.g., IXO 1, 2 in

NGC 720; see Table 1) the degree of contamination by foreground/background objects is very important. In any case, the contamination is not total, so it is incorrect to think that all ULXs are QSOs with different redshifts (Burbidge et al. 2003).

Several statistical analyses (e.g. Ptak & Colbert 2004; Schwartz et al. 2004; Irwin et al. 2004) of current compilations of ULXs confirm the relevance of such contamination. In this paper we address the problem of such a contamination in existing samples of ULXs, extending the work by these authors. We analyse the catalogue by Colbert & Ptak (2002) of ULX candidates and estimate the contamination in each galaxy and the most likely truly ULX sources. The paper is organized as follows: Section 2 describes the catalogue used and the corrections applied to obtain an homogeneous sample; the distribution of background sources and the method of computing probabilities are considered in Sections 3 and 4; Section 5 presents the results of the statistical analysis; and Sections 6 and 7 present the discussion and conclusions respectively, analysing also the possibility of some non-cosmological redshifts.

2. Catalogues of ULXs and sample selection

There are several catalogues of ULXs (ultraluminous X-ray sources), also called IXOs (intermediate-luminosity X-ray objects), built from ROSAT, XMM or Chandra data. Colbert & Ptak (2002) (C02 hereafter), and Liu & Bregman (2005) are based in ROSAT data, while Schwartz et al. (2004) used Chandra data, and Liu & Mirabel (2005) did a compilation of

the literature with a mixture of different data from different surveys, including data from Chandra and XMM X-ray telescopes. We decided to analyse the C02 catalogue because i) an important fraction of the sources have been identified in the optical, and this allows a direct comparison with our statistical results; ii) the objects have been selected using data from just one X-ray instrument, doing a homogeneous sample; and iii) a correction is possible that includes objects that were known to be background objects.

The analysis should not be carried out on the sample as a whole because we are dealing with a catalogue with constraints in luminosity rather than in flux for galaxies within a wide range of distances; we cannot do a simple calculation such as multiplying the total covered area by the average density of QSOs up to a given magnitude, because the limiting magnitude is different for each galaxy. Analysing the galaxies individually will also enable us to identify the galaxies likely to be totally/partially contaminated and those that are not, and will provide a useful tool for ULX searches and optimizing future follow-up studies. Furthermore, we compile the recent literature about the observation of the optical counterparts of these candidate ULXs that will corroborate our statistical analysis or will show whether there is some possible anomalous excess of different redshift objects around the parent galaxies.

C02 presented a catalogue of 87 candidate ULXs, which were observed in X-ray with ROSAT HRI (Voges et al. 1999); they were around bright galaxies from the RC3 catalog whose velocity is less than 5000 km/s and have more than 90% of the galaxy in the field of view of the ROSAT HRI observations (the total number of galaxies within these constraints was 766). They were detected by simply measuring the luminosity of these ROSAT sources assuming they are at the distance of a parent galaxy that is nearby: their sample of IXOs follows the constraint that the ULX is at an angular distance less than $2R_{25}$ from the centre of an RC3 galaxy, where R_{25} is the major axis of the 25 mag/arcsec² isophote, and the luminosity assuming the ULX has the same distance as the galaxy $L_X[2-10keV] \geq 10^{39}$ erg/s. A total of 54 galaxies had at least one of the 87 candidate ULXs with these constraints, and some galaxies were surrounded by up to seven candidate ULXs.

C02 have not included in their catalogue some sources observed by ROSAT HRI that could be classified as candidate ULXs according to their constraints because they were known to be QSOs with a redshift different from that of the RC3 galaxy. Arp et al. (2004, hereafter A04) (table 1) list these sources; there are 12 extra candidate ULXs in this list which follow the C02 constraints. Even if they are background sources that have been discarded as ULXs associated with the galaxy, they should be included in the catalogue if we wish to select a statistical sample. Indeed, as we will see throughout this paper, many of the candidate ULXs turned out later also to be background sources, so there is no reason to separate sources that were known to be background sources before or after the C02 compilation. If we include these sources, there are 99 candidate ULXs around 57 galaxies. Hereafter we will denote this sample as C02c ('Colbert & Ptak 2002 corrected'). They are listed in Table 1.

3. Distribution of background sources

To estimate the contamination in the C02c catalogue we consider the distribution of background sources in two different wavelength ranges: i) any source, $W \equiv X$, X-ray at frequencies between 0.5 and 2.0 keV; ii) QSOs, $W \equiv B$, B optical filter.

For the X-ray, we use the cumulative counts by Hasinger et al. (1993) obtained in the Lockman Hole from ROSAT-PSPC counts using the flux spectral-shape they also derived. A parameterization of their fig. 8 with two linear regimes gives (in units of deg⁻²) as a function of the flux $S_{X,i}$ (units: erg/s/cm²):

$$N_{X,i>(> S_i) = \begin{cases} 1.0 \times 10^{-21} S_{X,i}^{-1.66}, & S_{X,i} \geq 2 \times 10^{-14} \\ 2.8 \times 10^{-12} S_{X,i}^{-0.97}, & S_{X,i} < 2 \times 10^{-14} \end{cases} \quad (1)$$

This law is valid in principle for the range $10^{-15} < S_{X,i} < 10^{-12}$, where most of our ROSAT sources are. Hasinger (1998) gives the counts for a wider regime, and the linear regimes are present in wider ranges, although the exponents change slightly for larger values of $S_{X,i}$. Zickgraf et al. (2003, fig. 18) give the counts for brighter X-ray sources. In any case, for the estimation of probabilities (see next section), the law of eq. (1) is accurate enough.

We take the ROSAT X-ray flux of the sources, $S_{X,i}$, from C02c. They give the luminosity for the range $2 \text{ KeV} < \nu < 10 \text{ KeV}$ but this can be converted into fluxes if we divide the luminosity by $4\pi d_{galaxy}^2$; and $S_{X,i} \equiv S_i[0.5 \text{ KeV} < \nu < 2 \text{ KeV}] = 0.548 S_i[2 \text{ KeV} < \nu < 10 \text{ KeV}]$. The factor 0.548 stems from the assumption that the X-ray spectrum of the sources is $S[\nu]d\nu \propto \nu^{-0.7}d\nu$, an assumption also adopted by C02 to calculate their luminosities.

For the estimation of the background QSOs in the B-band, we use the counts by Cheney & Rowan-Robinson (1981) for $m_B \leq 18.5$ and Boyle et al. (2000) for $m_B > 18.5$ (see the fit in fig. A.1 of López-Corredoira & Gutiérrez 2004) as a function of the photographic magnitude $m_{b,i}$. Where no B-magnitude is available for a candidate ULX optical counterpart, we use the V-magnitude (in QSOs, $(B - V)$ is small). Since we are only interested in a rough approximation of the counts, no colour correction for the difference between photographic and Johnson filter or with V in case of few QSOs from A04 is applied. Boyle et al. (2000) counts of colour-preselected QSOs agree more or less with the counts from a more complete spectroscopic survey by Meyer et al. (2001). The counts include both QSOs and Seyfert 1 galaxies (Seyfert 1-type galaxies would form ~10% of the total counts) but not Seyfert 2, LINERs or other types of AGNs. According to Hao et al. (2005), Seyfert 2 would be less than four time more numerous than Seyfert 1, so the inclusion of Seyfert 2 would increase the counts up to 40%. Most of our sources are presumably point-like QSOs rather than extended galaxies, given the images of the optical counterparts. In any case, since we are interested in the order of magnitude, it is not so important whether an AGN is a QSO or a Seyfert galaxy.

The counts are (units deg⁻²):

$$N_{B,i(< m_{B,i})} = \begin{cases} 10^{-2.8+0.8(m_{B,i}-15)}, & m_{B,i} \leq 18.5 \\ 1981.0 - 214.2m_{b,i} + 5.792m_{b,i}^2, & m_{B,i} > 18.5 \end{cases} \quad (2)$$

In this case, the B magnitudes were taken from the NOMAD database¹ of the most likely optical counterpart (the brightest candidate closest to and within 10'' of the X-ray source). The nominal positional accuracy of ROSAT HRI is 5 arcseconds (Voges et al. 1999). If we compare the position of the 14 candidate ULXs common to CPO2 (with ROSAT) and Schwartz et al. (2004) (with Chandra), we find that the average difference is 2.4 arcseconds. If we do not find any counterpart within 10'', we attribute this a magnitude $m_B > 21.5$, i.e. higher than the limiting magnitude of the Palomar plates. It may happen that the optical source is the false counterpart because the real ULX is much fainter; however, the probability is not very high in regions away from the galaxy. In fact, the probability of observing an object with magnitude ~ 20 within a circle of radius 10'' is $\leq 10^{-3}$ for QSOs (Boyle et al. 2000) and $\sim 10^{-2}$ for high galactic latitude stars (Robin et al. 2003), which we consider negligible. A higher contamination could be present in spiral arms, where the density of new-born stars over HII regions could be high. For instance, Kaaret (2005) shows that IXO 5 is much fainter than the magnitude of the brightest source in A04, found as an optical counterpart. In this case, and when spectroscopy of the optical counterpart has not revealed a QSO or stellar nature, the contamination with regard the visible magnitude would be underestimated but never overestimated, so we will get a *minimum* of the ratio of contamination.

4. Probabilities of being background sources

From the catalogue of ULXs in Table 1 we want to calculate the probabilities that they are background objects (mainly background QSOs) instead of X-ray sources associated with their putative parent galaxies.

Given a galaxy with n ULXs around whose angular distances from its centre are α_i (deg) and background density $N_{W,i}$ (deg^{-2}) of sources brighter than the flux/“apparent magnitude” of this source in a wavelength range W , the probability that all of them are background sources is:

$$P_W \sim e^{-\lambda_{W:\max}} \left(\sum_{i=n}^{\infty} \frac{\lambda_{W:\max}^i}{i!} \right) \left(\prod_{j=1, j \neq i_{W:\max}}^n \frac{2\lambda_{W,j}}{\lambda_{W:\max}} \right), \quad (3)$$

$$\lambda_{W:\max} = \text{Maximum}[\lambda_{W,i}]; \quad i_{W:\max} / \lambda_{W,i_{\max}} = \lambda_{W:\max}$$

$$\lambda_{W,i} = \pi \alpha_i^2 N_{W,i}$$

The first two factors represent a Poissonian distribution for n background sources equivalent to $i_{W:\max}$ (the fainter and more distant source). The third factor, $\left(\prod_{j=1, j \neq i_{W:\max}}^n \frac{2\lambda_{W,j}}{\lambda_{W:\max}} \right)$, takes into account that the sources might be brighter or closer to the centre of the galaxy than normally expected in a random process. Randomly, one would expect that $\lambda_j / \lambda_{\max} \approx 1/2$ on average, so this third factor should be nearly unity; but in some cases, some sources are much brighter than $i_{W:\max}$ or much closer to the centre than $i_{W:\max}$, and in such cases the third factor would be much lower than one, reflecting the fact of this improbable difference in brightness or angular distance.

For example, in the case with three candidate ULXs with magnitudes 20, 16 and 15 respectively and distances around 1 arcminutes from the nuclei of the galaxy, we will calculate the Poissonian probability of having background QSOs within a circle of 1 arcminute with magnitude lower than 20 (with the first two factors), and the third factor will reduce this probability to reflect the fact that the QSOs with magnitudes 15 and 16 are much brighter (improbable) than those with magnitudes 19–20. As another example, imagine three QSOs of roughly the same magnitude 20 with distances to the galaxy 0.1, 0.2 and 3.0 arcminutes respectively. We will calculate the Poissonian probability to have background QSOs within a circle of 3 arcminutes with magnitude 20 (with the first two factors), and the third factor will reduce this probability to reflect that the QSOs with distances 0.1 and 0.2 arcminutes are much closer than expected randomly.

Table 1 gives $\lambda_{X,j}$ and $\lambda_{B,j}$ for each source, and the global probabilities P_X and P_B for each galaxy. The X-ray fluxes and the B-band magnitudes are already listed in this table. And, if there is an optical identification, this is indicated in the column 9.

5. Results

5.1. Overall contamination

A rough calculation with X-ray probabilities will give the total number of background sources:

$$\text{number of background sources} \approx \sum_{j=1}^{99} (1 - e^{-13.4\lambda_{X,i}}) = 50.3, \quad (4)$$

where the factor $13.4 = 766/57$ is the ratio of the total number of galaxies to those with at least one candidate ULX. The total number of background sources that we obtain with this expression is 51% of the background sources. It would be > 51.2 ($> 52\%$, only QSOs) if we did the calculation with $\lambda_{B,i}$ instead of $\lambda_{X,i}$.

So far, 54 sources have been already identified if we include the diffuse objects as identified (see Table 1), and 36 of them ($67 \pm 11\%$) turned out to be contamination; 18 of the ULX candidates are indeed sources belonging to the parent galaxy. This is slightly higher than our statistical estimate in X-ray. We must also bear in mind that there may be a bias in the identified sources. They are on average brighter in the visible than the unobserved sources, although in principle fainter objects should have an even higher probability in the visible to be contamination. Other selection effects such as the avoidance of crowded regions in the spiral arms near the centre of the galaxies, where many optical counterparts are possible, could reduce this probability.

Ptak & Colbert (2004) calculated a similar contamination: around 44%. Liu et al. (2005b) calculated a contamination ratio about a 30% within $2R_{25}$. Swartz et al. (2004) conclude that $\sim 25\%$ of the sources ULXs candidates from Chandra ACIS may be background objects including 14% of the ULX candidates in the sample of spiral galaxies and 44% of those in

¹ <http://www.navy.mil/nomad.html>

elliptical galaxies, but their numbers are not directly comparable with that of the C02 sample because in a fair number of cases the Chandra fields do not cover the circles within $2R_{25}$. For Irwin et al. (2004) the contamination is nearly total for elliptical galaxies and for the sources far from the centre in spiral galaxies. As said in the introduction, other authors (Kilgard et al. 2002; Schwartz et al. 2004; Liu et al. 2005b) also predicted a lower contribution of background sources for late-type galaxies.

5.2. Contamination in each galaxy

Table 1 presents the results of the statistical analysis. The expected mean number of random sources within the angular area are given in columns 4 and 7 for the analyses in the X-ray and in the optical respectively. The probabilities that all the ULXs detected in a given galaxy are contaminants are given in columns 5 and 8. Since C02c had a sample of 766 galaxies from which there are 57 galaxies with at least one candidate ULX, only galaxies with P_X or P_B lower than 7×10^{-5} will have within 95% C.L. at least one X-ray source belonging to the galaxy instead all of them of being background sources. Of the 57 galaxies, only 18 of them follow this: NGC 1073, NGC 1313, NGC 1399, NGC 1672, Holmberg II, NGC 3628, NGC 4088, NGC 4319, NGC 4374, NGC 4472, NGC 4485, NGC 4559, NGC 4631, NGC 4649, NGC 4861, NGC 5128, NGC 5204 and NGC 5194. The average Hubble type of these galaxies is +3.2. We found some ellipticals (Hubble type: -5 or -4) with a low probability of all their ULXs being background sources, although no ULX was confirmed spectroscopically in them. See §6 for a discussion on their low probabilities of being background sources.

The second group with the remaining 39 galaxies would have 53 candidate ULXs, which would be mostly background sources: either QSOs or any other kind of X-ray emitter. The average Hubble type of the galaxies of this second group is +2.0. Indeed 22 sources of this second group have already a spectroscopic identification and 19 of them are QSOs or emission line galaxies with different redshifts from that of the main galaxy (see Table 1). Another source is a foreground M star. Moreover, if one looks at the photometry of the fourth release of the SDSS (Adelman-McCarthy et al. 2006), one will find within $10''$ optical counterparts of seven IXOs of this second group (4, 44, 48, 63, 64, 74 and 83) and three of them (4, 44 and 74) have colours typical of QSOs [according to the criterion in eq. (1) of López-Corredoira & Gutiérrez 2004, which means approximately that they have a probability of being real QSOs of 2/3].

This statistical analysis implies that, if one looks for ULXs, one should preferably observe the candidate ULXs of the 18 galaxies from the first group (with P_X or P_B lower than 7×10^{-5}). For many of these 18 galaxies, spectroscopic identification is also available: there are confirmed sources belonging to the main galaxy for IXOs 5, 7, 8, 31, 65, 68, 72, 73, 77, 80, 81 (see Table 1). Some other candidate ULXs are known to be bright diffuse sources that cannot be background QSOs: IXOs 26, 27, 42, 62, 79. And there are also some identified QSOs

with redshifts different from those of the parent galaxies: IXOs 19, 20, 57, 58, 69, 71; A3, 5, 9, 10, 13, 14, 21, 22. IXO75 is a foreground local M star; therefore, there remain still to be identified 15 IXOs of the first group: IXOs 17, 18, 25, 39, 50, 51, 55, 56, 59, 60, 61, 66, 70, 76, 78. The variable X-ray source IXO 76 has a possible optical counterpart corresponding to a late O or early B star with magnitude 24.1 in the F555W filter (Ghosh et al. 2005), but this has not been confirmed spectroscopically. If one looks at the fourth release of the SDSS (Adelman-McCarthy et al. 2006), one will find optical counterparts of IXOs 50, 51, 55, 56, 59, 60, 61 and 70, among which IXOs 50, 51, 55, 59, 70 have colours typical of QSOs [according to the criterion in eq. (1) of López-Corredoira & Gutiérrez 2004; although they might also be HII regions with the same colours], while IXOs 56, 60, 61 have a different colour, so they should be ULXs, foreground stars, or emission line galaxies.

5.3. Statistics with the clean sample

We define the clean sample as the objects that are confirmed ULXs or candidates with low probabilities of being contaminants.

According to the discussion in the previous subsection, the objects that have not yet been observed spectroscopically, and that have higher probabilities of being ULXs in the C02 catalogue are IXOs 17, 18, 25, 39, 66, 76 and 78. We removed IXOs 50, 51, 55, 59 and 70 from the initial list for their typical QSOs colours without being diffuse nebulae. We also removed IXOs 56, 60, 61 because their $\lambda_{B,j} > 1$ while the P_B and P_X of the parent galaxy NGC 4472 are not so low as to forbid a projection of seven QSOs (or any other kind of contaminating object).

The objects that are spectroscopically confirmed ULXs are (see Table 1) IXOs 5, 7, 8, 22, 31, 34, 65, 68, 72, 73, 77, 80, 81; plus IXOs 26, 27, 42, 62, 79, which were not observed spectroscopically but are bright diffuse sources, so they cannot be background QSOs, and are most probably HII regions of the parent galaxies.

In total, there are 25 objects. This sample might not be totally clean; even the objects identified as HII of the parent galaxy might correspond to wrong optical counterparts, but at least the contamination will be much lower than the original C02 sample. We will repeat statistics made by Ptak & Colbert (2004) on C02 sample, but only with these 25 objects. A histogram of energy distribution is shown in Figure 1, and a histogram of the distribution according to Hubble type is presented in Figure 2.

The distribution of energy peaks in intermediate energies ($\sim 4 \times 10^{39}$ erg/s), or could be flat for lower energies (the Poissonian errors do not allow a distinction). The distribution of the Hubble types peaks around late-type spirals, as expected; however, we have still two ellipticals. An examination of the IXOs from these ellipticals makes us realize that they come from the unobserved candidates around the elliptical galaxies NGC 1399 and NGC 5128. One is tempted to claim that these candidate ULXs are indeed background QSOs, but in such a case, we would have the problem of an anomalously high con-




Fig. 1. Distribution of luminosities in the clean sample of ULXs.




Fig. 2. Distribution of Hubble types of the galaxies with at least one member of the clean sample of ULXs.

centration of QSOs around NGC 1399 (see §6) and an inexplicably low value of P_X in NGC 5128 due to contamination. NGC 1399 is indeed a cD galaxy at the centre of a cluster; perhaps the ULXs near NGC 1399 belong to nearby galaxies in the cluster. The three ULXs in these two galaxies (IXOs 17, 18, 76) have not yet been observed; they are very faint (no optical counterpart has yet been found), so we cannot confirm whether the expectation by Irwin et al. (2004) that there are no ULXs associated with ellipticals is correct or not.

6. QSOs with low probability to be background sources

ULXs around some galaxies have been proposed (Burbidge et al. 2003) to be QSOs or BL Lac objects with intrinsic redshift being ejected from a parent galaxy rather than black holes with

masses $10^2 - 10^4 M_\odot$. It is clear, as said in the introduction, that this cannot be the case for all candidate ULXs because there are many whose spectroscopy have shown a different nature. We have tested whether the Burbidge et al. hypothesis could be applied to some of the objects. Let us do a statistical analysis for each individual galaxy of the first group (with P_X or P_B lower than 7×10^{-5}) in which some QSO has been observed:

NGC 1073 : with A3,5 as QSOs. Since IXO 5 is not a QSO, the probability of being background objects is not so low for only two QSOs (excluding IXO 5 from the list): $P_X = 10^{-3}$, $P_B = 10^{-4}$; quite possibly background sources. Note, however, that there is a third QSO around NGC 1073 which was not included here because the X-ray flux is less than the C02 requirements (equivalent to a luminosity $L_X[2 - 10keV] \geq 10^{39}$ erg/s if they were at the distance of the galaxy). With this third QSO, the probability becomes lower, as claimed by Arp & Sulentic (1979).

NGC 1399 : with IXOs 19, 20 as QSOs. Were IXOs 17, 18 also QSOs, the probability of their being background sources would be very low; however, their optical spectra have still not been obtained, and they will probably not be observed in the near future because they are quite faint (they have no optical counterpart in the Palomar plates). For the two discovered QSOs (excluding IXOs 17, 18 from the list) the probabilities are: $P_X = 3 \times 10^{-3}$, $P_B = 10^{-5}$. We suspect that there may be an error in the magnitudes given by NOMAD for the optical counterpart of IXO 20 ($m_B = 15.2$, $m_R = 18.1$).

NGC 3628 : with A9,10,13,14 as QSOs. Again, the most intriguing case is IXO 39, with lower values of λ , but this has not been observed spectroscopically yet since its optical counterpart is not detected. With the remaining four QSOs (excluding IXO 39 from the list): $P_X = 2 \times 10^{-5}$, $P_B = 8 \times 10^{-5}$. And there are still another three QSOs in A04 which do not follow the constraint $L_X[2 - 10keV] \geq 10^{39}$ but which, if included, will considerably reduce the probability to three orders of magnitude. Precisely because of that, it is a known anomalous redshift candidate (Arp 2002), in which there is not only the overdensity but also an alignment of QSOs across the nucleus of NGC 3628 and their connection with X-ray isophotes and coincidences in the alignment with some other filaments in the radio and optical.

NGC 4319 : with A21 as QSO. This known QSO, Mrk 205, is indeed a well known controversial case in the debate about anomalous redshifts. It is not only the low probability of finding so bright a QSO near NGC 4319 ($P_X = 4 \times 10^{-6}$ and $P_B = 10^{-6}$) but also the fact that it is connected by a small bridge to the galaxy (Sulentic & Arp 1987a,b). There is also an almost continuous luminous connection extending from Mrk 205 into the nucleus of the spiral, and a corresponding feature on the opposite side of the disc appearing to link a bright UV knot ($m_B = 19.5$; with redshift of the galaxy, Sulentic & Arp 1987b) with the nucleus (Sulentic & Arp 1987a).

NGC 4374 : with A22 as QSO. The probabilities alone for this object (excluding IXOs 50 and 51 from the list) are:

$P_X = 2 \times 10^{-2}$, $P_B = 3 \times 10^{-3}$, high enough for A22 most likely to be a background object. IXOs 50 and 51 have not yet been observed. There is no optical counterpart in the Palomar plates; however there are optical counterparts in SDSS-4th release survey: $u = 24.63$, $g = 25.11$, $r = 24.80$ for IXO 50 (with a probable large error in these magnitudes because we are at the limit of the detection) and $u = 20.97$, $g = 20.70$, $r = 20.01$ for IXO 51. Both of them might be QSOs according to their colour [see eq. (1) of López-Corredoira & Gutiérrez 2004]. Were these two cases really QSOs, we would have a very interesting new case of anomalous redshift.

NGC 4472 : with IXOs 57, 58 as QSOs. And there remain IXOs 55, 56, 59, 60 and 61 to be identified. Only with the IXOs 57 and 58 as QSOs (excluding the other five IXOs from the list), the probabilities that they are both background objects are: $P_X = 0.1$, $P_B = 2 \times 10^{-3}$. They are most probably two background QSOs. IXOs 55 ($u = 20.57$, $g = 20.66$, $r = 20.41$) and 59 ($u = 20.68$, $g = 20.07$, $r = 20.08$) have SDSS colours of QSOs [see eq. (1) of López-Corredoira & Gutiérrez 2004], while IXOs 56 ($u = 21.53$, $g = 20.71$, $r = 20.59$), 60 ($u = 22.89$, $g = 21.36$, $r = 20.81$), and 61 ($u = 23.42$, $g = 22.22$, $r = 21.35$) are redder. Even where we had four QSOs (IXOs 55, 57, 58 and 59), the probability would be high enough to be a fortuitous case of background sources.

NGC 4649 : with IXOs 69 and 71 as QSOs. IXO 70 remains to be identified, although according to its SDSS colour ($u = 19.34$, $g = 19.32$, $r = 19.09$), it might also be a QSO. Only with the IXOs 69 as 71 as QSOs (excluding IXO 70 from the list), the probabilities that they are both background objects are: $P_X = 10^{-2}$, $P_B = 2 \times 10^{-3}$. They are most probably two background QSOs.

Therefore, the possibly controversial cases with regard to anomalous redshift are only those that were known in the past, such as NGC 3628 and NGC 4319; the C02 catalog does not contain clear cases of QSOs with different redshifts and low probabilities of being background objects.

Radecke (1997) points out a detection of correlation between nearby Seyfert galaxies and X-ray sources, but the selection is different from those used in the C02 catalogue. Radecke uses only Seyfert galaxies while we use all kinds of galaxies. Radecke gets the maximum correlation for very bright sources with B magnitude lower than 10.5, while most galaxies in C02 sample are fainter. Finally, Radecke puts the constraint for X-ray sources having more than 5 counts/ks while in the C02 sample most of the sources have less than 5 counts/ks. Therefore, there is no inconsistency between Radecke's analysis (which finds a number of X-ray sources around his galaxies considerably higher than expected from background sources; in the best of the cases, he gets excess over 250%) and our result (we got an excess of $\sim 100\%$ over the background sources, which are due to ULXs belonging to the galaxies but they are not QSOs). It is also quite possible that part or all of the excess of sources around Seyfert galaxies in Radecke's analysis is due to the existence a considerable number of ULXs belonging to the galaxies, these ULXs having the same redshift as the galax-

ies, so they could not be used as a claim of anomalous redshift. However, this hypothesis has not been tested yet.

7. Conclusions

From the above analysis we conclude the following:

- More than a half of the ULX candidate in the C02c catalogue are background QSOs. This result from the statistical analysis is confirmed by follow-up spectroscopy of some of the sources.
- The clean sample is as follows. The spectroscopically confirmed ULXs are IXOs 5, 7, 8, 22, 31, 34, 65, 68, 72, 73, 77, 80, 81; plus IXOs 26, 27, 42, 62, 79, which were not observed spectroscopically but are bright diffuse sources, so they cannot be background QSOs and are most probably HII regions of the parent galaxies. Other objects that have not yet been observed spectroscopically, and that have higher probabilities of being ULXs in the C02 catalogue are IXOs 17, 18, 25, 39, 66, 76 and 78.
- Among the 87 sources in C02 catalogue, there are no significant statistical cases that indicate non-cosmological redshifts in some QSOs associated with galaxies. Nonetheless, the problem of the anomalous redshifts is still open because there remain other known cases, for instance those of NGC 4319 or NGC 3628, with significantly low probabilities of their QSOs being background objects.

Acknowledgments: Thanks are given to the referee, P. Kaaret, for helpful comments. The authors were supported by the *Ramón y Cajal* Programme of the Spanish Science Ministry. Thanks are given to T. J. Mahoney (IAC) for the proof-reading of the paper.

References

- Adelman-McCarthy, J. K., Agüeros, M. A., Allam, S. S., et al., 2006, *ApJS* 162, 38
- Arp, H. C., 1977, *ApJ* 218, 72
- Arp, H. C., 2002, *A&A* 391, 833
- Arp, H. C., Gutiérrez, C. M., López-Corredoira, M., 2004, *A&A*, 418, 877 (A04)
- Arp, H. C., & Sulentic, J. W. 1979, *ApJ* 229, 496
- Boyle, B. J., Shanks, T., Croom, S. M., et al., 2000, *MNRAS*, 317, 1014
- Burbidge G., Burbidge E. M., & Arp H., 2003, *A&A* 400, L17
- Cheney, J. E., & Rowan-Robinson, M., 1981, *MNRAS*, 195, 497
- Clark, D. M., Christopher, M. H., Eikenberry, S. S., et al., 2005, *ApJ* 631, L109
- Colbert, E. J. M., & Mushotzky, R. F., 1999, *ApJ*, 519, 89
- Colbert, E. J. M., & Ptak, A. F., 2002, *ApJS*, 143, 25 (C02)
- Gao, Y., Wang, Q. D., Appleton, P. N., & Lucas, R. A., 2003, *ApJ* 596, L171
- Ghosh, K. K., Finger, M. K., Swartz, D. A., Tennant, A. F., & Wu, K., 2005, *astro-ph/0509720*
- Ghosh, K. K., Swartz, D. A., Tennant, A. F., Saripalli, L., Gandhi, P., Foellmi, C., Gutiérrez, C. M., & López-Corredoira, M., 2006, in preparation
- Gutiérrez, C. M., 2006, *ApJ-Lett.*, accepted; *astro-ph/0602207*

- Gutiérrez, C. M., López-Corredoira, M., 2005, ApJ, 622, L89
- Gutiérrez, C. M., López-Corredoira, M., 2006, in preparation
- Hao, L., Strauss, M. A., Fan, X., et al. 2005, AJ, 129, 1795
- Hasinger, G., 1998, Astron. Nachr., 319, 37
- Hasinger, G., Burg, R., Giacconi, R., Hartner, G., Schmidt, M., Trümper, J., & Zamorani, G., 1993, A&A, 275, 1
- Irwin, J. A., Bregman, J. N., & Athey, A. E., 2004, ApJ 601, L143
- Kaaret, P., 2005, ApJ 629, 233
- Kaaret, P., Ward, M.-J., & Zezas, A. 2004, MNRAS 351, L83
- Kilgard, R. E., Kaaret, P., Krauss, M. I., Prestwich, A. H., Raley, M. T., & Zezas, A., 2002, ApJ, 573, 138
- Larsen J. A., Humphreys R. M. 2003, AJ 125, 1958
- Lehmann, I., Becker, T., Fabrika, S., et al. 2005, A&A 431, 847
- Lira, P., Lawrence, A., & Johnson, R. A., 2000, MNRAS, 319, 17
- Liu, J.-F., Bregman, J.-N., 2005, ApJS 157, 59
- Liu, J.-F., Bregman, J.-N., & Seitzer, P., 2004, ApJ 602, 249
- Liu, J.-F., Bregman, J.-N., Seitzer, P., & Irwin, J., 2005a, astro-ph/0501310
- Liu, J.-F., Bregman, J. N., & Irwin, J., 2005b, astro-ph/0501312
- Liu, Q.-Z., & Mirabel, I. F., 2005, A&A 429, 1125
- López-Corredoira, M., & Gutiérrez, C. M., 2004, A&A, 421, 407
- Makishima, K., Kubota, A., Mizuno, T., et al., 2000, ApJ 535, 632
- Meyer, M. J., Drinkwater, M. J., Phillips, S., & Couch, W. J., 2001, MNRAS, 324, 343
- Mucciarelli, P., Zampieri, L., Falomo, R., Turolla, R., & Treves, A., 2005, ApJ 633, L101
- Pakull, M. W., & Mirioni, L., in: New Visions of the X-ray Universe in the XMM-Newton and Chandra Era, ESTEC, The Netherlands. Preprint astro-ph/0202488
- Ptak, A., & Colbert, E., 2004, ApJ 606, 291
- Radecke, H. D., 1997, A&A, 319, 18
- Ramsey, C. J., Williams, R. M., Gruendl, R. A., Chen, C.-H. R., Chu, Y.-H., & Wang, Q. D., 2005, astro-ph/0511540
- Robin, A. C., Reylé, C., Derrière, S., & Picaud, S., 2003, A&A, 409, 523
- Smith, B. J., Struck, C., Nowak, M. A. 2005, AJ 129, 1350
- Sulentic, J. W., & Arp, H. C. 1987a, ApJ 319, 687
- Sulentic, J. W., & Arp, H. C. 1987b, ApJ 319, 693
- Swartz, D., Ghosh, K., Tennant, A., & Wu, K., 2004, ApJS 154, 519
- Voges, W., et al. 1999, in Diffuse thermal and relativistic plasma in galaxy clusters, ed. H. Bohringer, L. Feretti, & P. Schuecker (Garching: Max-Planck-Institut für Extraterrestrische Physik), 179
- Wong D. S., Chornock R., & Filippenko A. V. 2005, *Populations of High Energy Sources in Galaxies, Proceedings of IAU Symposium No. 230, Eds. Evert J.A. Meurs & Giuseppina Fabbiano*, in press
- Zampieri, L., Mucciarelli, P., Falomo, R., Kaaret, P., Di Stefano, R., Turolla, R., Chierigato, M., & Treves, A., 2004, ApJ, 603, 523
- Zickgraf, F.-J., Engels, D., Hagen, H.-J., Reimers, D., & Voges, W., 2003, A&A 406, 535

Table 1. Column 1: RC3 galaxy. Column 2: number of IXO in the C02 catalog, or from table 1 in A04 (named with Axx, and the number xx of row in that table). Column 3: flux in the range from 0.5 to 2.0 keV [units 10^{-14} erg/s/cm²]. Column 4: average number of expected sources up to the distance of this source to the centre of the galaxy, and brighter than its flux in X-ray. Column 5: probability that all candidate ULXs associated to the galaxy are indeed background sources with the observed distribution of X-ray fluxes. Column 6: B magnitude of the most likely optical counterpart from NOMAD database; letter “d” indicates that it is a diffuse source or a point-like source embedded in a halo; letter “a” indicates that the source is in the spiral arm of the galaxy. Column 7: average number of expected sources up to the distance of this source to the centre of the galaxy and brighter than its B magnitude. Column 8: probability that all candidate ULXs associated with the galaxy are indeed background sources with the observed distribution of B magnitudes. Column 9: spectroscopic identification [reference]. References of the spectroscopic identification: 1: A04; 2: Zampieri et al. (2004); 3: Gutiérrez & López-Corredoira (2006); 4: Gutiérrez (2006); 5: Lehmann et al. (2005); 6: Kaaret et al. (2004); 7: Gutiérrez & López-Corredoira (2005); 8: Ramsey et al. (2005); 9: Liu et al. (2005a); 10: NED database; 11: Liu et al. (2004); 12: Arp (1977); 13: Wong et al. (2005) [We call ‘QSO’ those objects classified by Wong et al. (2005) as AGN, since most of these sources are presumably point-like QSOs rather than extended galaxies]; 14: Ghosh et al. (2006); 15: Pakull & Mirioni (2002); 16: Mucciarelli et al. (2005); 17: Kaaret (2005).

Galaxy	IXO	$S_{X,i}$	$\lambda_{X,i}$	P_X	$m_{B,i}$	$\lambda_{B,i}$	P_B	spectr. ident.
NGC 720	1	1.1	0.97	5×10^{-2}	20.1	0.16	7×10^{-4}	QSO [1,13]
	2	4.4	8.9×10^{-2}		18.7	5.1×10^{-3}		QSO [1,13]
NGC 891	3	7.9	0.16	0.1	> 21.5	> 1.46	> 0.8	
NGC 1042	4	17	3.5×10^{-3}	3×10^{-3}	> 21.5(a)	> 0.12	> 0.1	
NGC 1073	5	2.2	0.12	10^{-4}	13.1(a)	1.2×10^{-7}	8×10^{-12}	HII [17]
	A3	2.8	5.5×10^{-2}		V=20.0	2.4×10^{-2}		QSO [1]
	A5	7.0	2.7×10^{-2}		V=18.8	4.5×10^{-3}		QSO [1]
NGC 1291	6	9.8	5.8×10^{-2}	6×10^{-2}	> 21.5	> 0.78	> 0.5	
NGC 1313	7	210	1.5×10^{-5}	6×10^{-8}	> 21.5	> 3.2×10^{-2}	> 6×10^{-3}	HII [15]
	8	84	3.8×10^{-3}		20.0	0.47		O-BV or GI in HII [2,8,15,16]
NGC 1316	9	8.0	0.59	2×10^{-3}	19.5	0.70	> 3×10^{-4}	QSO [13]
	10	4.0	0.27		20.2	0.27		QSO [13]
	11	2.5	0.36		> 21.5	> 0.51		
	12	2.0	1.04		> 21.5	> 1.01		
	13	2.0	2.95		21.2	2.33		
	14	2.5	3.05		19.1	0.23		QSO [13]
	15	5.9	0.29	4×10^{-2}	19.4	0.17	10^{-3}	QSO [13]
NGC 1365	16	5.9	1.01		18.7	2.7×10^{-2}		QSO [13]
	A6	5.9	0.48		V = 19.7	0.47		QSO [1]
	17	3.1	2.9×10^{-2}	10^{-7}	> 21.5	> 5.7×10^{-2}	> 6×10^{-9}	
NGC 1399	18	6.2	4.2×10^{-3}		> 21.5	> 2.6×10^{-2}		
	19	31	1.4×10^{-2}		20.1	0.37		QSO [13]
	20	4.9	0.23		15.2	4.3×10^{-5}		QSO [13]
	21	5.0	4.3×10^{-3}	4×10^{-3}	18.4	2.9×10^{-4}	3×10^{-4}	
IC342	22	48	6.5×10^{-3}	6×10^{-3}	17.8	6.3×10^{-3}	6×10^{-3}	HII [3,8,15]
NGC 1553	23	5.5	0.10	0.1	20.8	0.32	0.3	
NGC 1566	24	5.8	4.8×10^{-3}	5×10^{-3}	> 21.5(a)	> 2.7×10^{-2}	> 3×10^{-2}	
NGC 1672	25	1.8	0.11	2×10^{-5}	> 21.5(a)	> 9.8×10^{-2}	$\sim 2 \times 10^{-14}$	
	26	5.6	1.1×10^{-2}		15.1(da)	2.9×10^{-4}		
	27	4.5	2.5×10^{-2}		R=13.4(da)	$\sim 1.4 \times 10^{-7}$		
NGC 1792	28	3.4	1.3×10^{-2}	10^{-2}	> 21.5	> 3.0×10^{-2}	> 3×10^{-2}	
NGC 1961	29	5.3	7.2×10^{-2}	7×10^{-2}	20.6	0.17	0.2	
Mrk 3	30	4.0	4.2×10^{-2}	4×10^{-2}	> 21.5	0.13	> 0.1	
Holmberg II	31	360	3.7×10^{-5}	4×10^{-5}	14.2	1.3×10^{-6}	10^{-6}	O4V-B3Ib in HII [5,6,8,15]
NGC 2775	32	2.2	0.55	6×10^{-2}	18.1	5.7×10^{-3}	2×10^{-4}	QSO [4,13]
	33	4.4	0.16		19.2	3.9×10^{-2}		QSO [7,13]
NGC 3031	34	360	1.4×10^{-3}	10^{-3}	20.4	3.00	0.95	4-6 Myr blue stars in HII [8,15]
NGC 3226	35	2.3	0.22	0.2	18.4	4.3×10^{-3}	4×10^{-3}	local M star [4,13]
NGC 3256	36	2.1	0.34	0.3	> 21.5	> 0.36	> 0.3	
IC 2597	37	23	3.0×10^{-3}	3×10^{-3}	20.8	9.5×10^{-2}	0.1	Emis.line Gal. [4,13]
NGC 3310	38	4.1	1.6×10^{-2}	2×10^{-2}	17.5	1.5×10^{-4}	10^{-4}	

Cont. table 1.

Galaxy	IXO	$S_{X,i}$	$\lambda_{X,i}$	P_X	$m_{B,i}$	$\lambda_{B,i}$	P_B	spectr. ident.	
NGC 3628	39	24	3.8×10^{-5}	3×10^{-10}	> 21.5	$> 2.3 \times 10^{-3}$	$> 8 \times 10^{-8}$		
	A9	770	9.0×10^{-2}		$V = 19.5$	0.10		QSO [1]	
	A10	15	4.9×10^{-2}		$V = 19.2$	9.3×10^{-2}		QSO [1]	
	A13	9.7	0.22		$V = 19.6$	0.42		QSO [1]	
	A14	31	7.8×10^{-2}		$V = 18.3$	9.1×10^{-2}		QSO [1]	
NGC 3923	40	7.3	0.11	0.1	19.0	3.7×10^{-2}	4×10^{-2}	QSO [4,13]	
UGC 7009	41	8.2	1.4×10^{-3}	10^{-3}	> 21.5	1.4×10^{-2}	10^{-2}		
NGC 4088	42	5.0	2.7×10^{-3}	3×10^{-3}	14.8(da)	2.4×10^{-7}	3×10^{-7}		
NGC 4151	43	2.2	1.02	6×10^{-4}	18.8	2.6×10^{-2}	8×10^{-4}	Emis.line Gal. [3,12,13]	
	44	1.1	0.46		20.9	0.16			
	A15	62	3.8×10^{-3}		$V = 20.3$	0.41		QSO [1]	
NGC 4168	A16	1.2	4.2×10^{-2}	4×10^{-2}	$V = 18.7$	4.2×10^{-4}	4×10^{-4}	QSO [1]	
NGC 4203	45	2.2	0.29	0.3	18.3	4.3×10^{-3}	4×10^{-3}	QSO [7]	
NGC 4254	46	14	6.2×10^{-3}	6×10^{-3}	> 21.5	> 0.15	> 0.1		
NGC 4319	A21	360	4.3×10^{-6}	4×10^{-6}	$V = 15.2$	9.9×10^{-7}	10^{-6}	QSO [1]	
NGC 4321	47	3.2	0.81	0.2	20.3	0.64	> 0.1		
	48	1.6	2.41		> 21.5	> 1.92			
	49	2.0	0.67		> 21.5	> 0.66			
	50	16	2.0×10^{-4}	2×10^{-7}	> 21.5	$> 6.3 \times 10^{-3}$	$> 6 \times 10^{-6}$		
NGC 4374	51	6.4	0.11		> 21.5	> 0.73			
	A22	10	1.8×10^{-2}		$V = 18.5$	3.1×10^{-3}		QSO [1]	
	52	2.2	0.20	0.2	18.7	3.7×10^{-3}	4×10^{-3}	QSO [13]	
NGC 4373	52	2.2	0.20	0.2	18.7	3.7×10^{-3}	4×10^{-3}	QSO [13]	
NGC 4395	53	35	3.1×10^{-3}	3×10^{-3}	> 21.5	> 0.36	> 0.3		
NGC 4438	54	13	2.6×10^{-2}	3×10^{-2}	19.5	6.9×10^{-2}	7×10^{-2}	QSO [3]	
NGC 4472	55	2.0	3.44	10^{-4}	20.8	2.03	$> 3 \times 10^{-5}$		
	56	4.1	1.14		20.8	2.13			
	57	10	0.21		18.2	3.4×10^{-2}		QSO [3]	
	58	4.1	0.96		18.4	4.7×10^{-2}		QSO [7]	
	59	3.2	2.19		21.0	3.27			
	60	13	9.6×10^{-2}		21.0	1.42			
	61	2.6	0.93		> 21.5	> 1.36			
	NGC 4485	62	11	2.7×10^{-4}	3×10^{-4}	13.2(d)	4.4×10^{-9}	4×10^{-9}	
	NGC 4552	63	5.1	0.26	5×10^{-2}	20.4	0.48	5×10^{-2}	
		64	3.2	0.25		20.0	0.14		
	NGC 4559	65	39	1.5×10^{-3}	5×10^{-8}	12.6	7.0×10^{-8}	$> 10^{-10}$	4 blue/red superg. in HII [3,9,15]
66		24	3.4×10^{-5}		> 21.5	$> 2.1 \times 10^{-3}$			
NGC 4565	67	21	5.0×10^{-4}	5×10^{-4}	> 21.5	$> 2.5 \times 10^{-2}$	$> 2 \times 10^{-2}$		
NGC 4631	68	12	1.6×10^{-2}	2×10^{-2}	$V=14.2(\text{da})$	$\sim 2.1 \times 10^{-6}$	$\sim 2 \times 10^{-6}$	HII [15]	
NGC 4649	69	5.2	8.8×10^{-2}	4×10^{-3}	18.3	5.4×10^{-3}	4×10^{-5}	QSO [7]	
	70	4.1	0.71		18.4	3.6×10^{-2}			
	71	10	0.16		19.8	0.46		QSO [7]	
	A23	10	4.8×10^{-3}	5×10^{-3}	$V = 20.5$	3.0×10^{-2}	3×10^{-2}	QSO [1]	
NGC 4861	72	9.1	4.9×10^{-3}	9×10^{-7}	9.5(a)	6.9×10^{-11}	$> 10^{-12}$	HII [10,15]	
	73	29	1.9×10^{-4}		> 21.5	$> 1.6 \times 10^{-2}$		HII [15]	
NGC 5055	74	35	1.3×10^{-2}	10^{-2}	18.4	2.3×10^{-2}	2×10^{-2}		
NGC 5128	75	38	9.2×10^{-3}	3×10^{-6}	10.7	1.3×10^{-8}	$> 4 \times 10^{-9}$	local M star [10,13]	
	76	150	2.9×10^{-4}		> 21.5	> 0.36			
NGC 5204	77	63	1.3×10^{-5}	10^{-5}	> 21.5	$> 3.8 \times 10^{-3}$	$> 4 \times 10^{-3}$	B0Ib in nebula [11,15]	
NGC 5194	78	6.5	3.8×10^{-2}	2×10^{-8}	> 21.5(a)	> 0.26	$> 5 \times 10^{-13}$		
	79	81	1.3×10^{-2}		13.0(da)	9.8×10^{-8}			
	80	16	8.1×10^{-3}		16.9(a)	2.5×10^{-4}		young star cluster [9]	
	81	13	1.4×10^{-2}		> 21.5(a)	> 0.29		? in young region [9]	
NGC 5236	82	33	9.8×10^{-3}	10^{-2}	> 21.5	> 0.98	> 0.6		
NGC 5457	83	20	0.12	0.1	> 21.5	> 4.99	> 0.99		
NGC 5775	84	2.3	0.43	0.3	19.6	7.6×10^{-2}	7×10^{-2}	QSO [7,14]	
NGC 6946	85	15	1.9×10^{-3}	2×10^{-3}	> 21.5(a)	$> 5.2 \times 10^{-2}$	$> 5 \times 10^{-2}$		
NGC 7314	86	3.2	6.4×10^{-2}	6×10^{-2}	18.6	1.7×10^{-3}	$> 2 \times 10^{-3}$		
NGC 7590	87	6.4	1.3×10^{-3}	10^{-3}	> 21.5(a)	$> 8.8 \times 10^{-3}$	$> 9 \times 10^{-3}$		